

NOTES ON NEW AND NOTEWORTHY PLANTS. XCIII

Harold N. Moldenke

AVICENNIA SCHAUERIANA f. GLABRESCENS Moldenke, f. nov.

Haec forma a forma typica speciei laminis foliorum ubique glabratis recedit.

This form differs from the typical form of the species in having both surfaces of its leaf-blades glabrate.

The type of the form was collected by Captain Reginald Charles Marshall (Herb. Trin. Bot. Gard. 12651) in the Caroni Swamp, Trinidad, on May 9, 1932, and is deposited in the Britton Herbarium at the New York Botanical Garden.

PAEPALANTHUS SESLERIOIDES var. CARABIAE Moldenke, var. nov.

Haec varietas a forma typica speciei statura minima foliis 1—3 cm. longis pedunculis 2.5—6.5 [—8] cm. longis recedit.

This variety differs from the typical form of the species in its much smaller stature, the leaves mostly only 1—3 cm. long and the peduncles usually only 2.5—6.5 cm. long, occasionally to 8 cm. long.

The type of the variety was collected by José Perez Carabia (no. 3941) on the Sabanas de los Indios, Isla de Pinos, Cuba, on May 11, 1940, and is deposited in the Britton Herbarium at the New York Botanical Garden.

PAEPALANTHUS SESLERIOIDES var. WILSONII Moldenke, var. nov.

Haec varietas a forma typica speciei statura perspicue parviora foliis 4.5—6.5 cm. longis pedunculis 8—20 cm. longis recedit.

This variety differs from the typical form of the species in its conspicuously smaller stature, the leaves mostly 4.5—6.5 cm. long and the peduncles mostly 8—20 cm. long.

The type of the variety was collected by Nathaniel Lord Britton, Elizabeth Gertrude Britton, and Percy Wilson (no. 14225) in white sand in the vicinity of Los Indios, Isla de Pinos, Cuba, on February 13, 1916, and is deposited in the Britton Herbarium at the New York Botanical Garden.

ADDITIONAL NOTES ON THE ERIOCAULACEAE. LXIV

Harold N. Moldenke

ERIOCAULON ODORATUM Dalz.

Additional synonymy: Eriocaulon odoratum Moldenke, Phytologia